

## 15-deoxy- $\Delta^{12,14}$ -PGJ<sub>2</sub>: endogenous PPAR $\gamma$ ligand or minor eicosanoid degradation product?

William S. Powell

*J Clin Invest.* 2003;112(6):828-830. <https://doi.org/10.1172/JCI19796>.

### Commentary

15-Deoxy- $\Delta^{12,14}$ -prostaglandin J<sub>2</sub> (15d-PGJ<sub>2</sub>) affects gene transcription by activating PPAR $\gamma$  and by covalent addition to transcription factors and signaling molecules. However, it is not known whether the high concentrations of 15d-PGJ<sub>2</sub> required for these responses are consistent with physiological levels. A new study suggests that in vivo 15d-PGJ<sub>2</sub> levels are actually several orders of magnitude below the levels required to induce many of the biological effects attributed to this molecule.

Find the latest version:

<https://jci.me/19796/pdf>



monitor and characterize CD4 T cell populations contributing to this disease. Such reagents may ultimately prove to be of use in monitoring the efficacy of intervention protocols that may inhibit progression to overt

IDDM in humans at risk for this debilitating disease.

1. Stratmann, T., et al. 2003. Susceptible MHC alleles, not background genes, select an autoimmune T cell reactivity. *J. Clin. Invest.* **112**:902-914. doi:10.1172/JCI200318337.

2. Lesage, S., et al. 2002. Failure to censor forbidden clones of CD4 T cells in autoimmune diabetes. *J. Exp. Med.* **196**:1175-1188.  
3. Verdaguer, J., Amrani, A., Anderson, B., Schmidt, D., and Santamaria, P. 1999. Two mechanisms for the non-MHC-linked resistance to spontaneous autoimmunity. *J. Immunol.* **162**:4614-4626.

## 15-deoxy- $\Delta^{12,14}$ -PGJ<sub>2</sub>: endogenous PPAR $\gamma$ ligand or minor eicosanoid degradation product?

William S. Powell

Meakins-Christie Laboratories, Department of Medicine, McGill University, Montreal, Quebec, Canada

15-Deoxy- $\Delta^{12,14}$ -prostaglandin J<sub>2</sub> (15d-PGJ<sub>2</sub>) affects gene transcription by activating PPAR $\gamma$  and by covalent addition to transcription factors and signaling molecules. However, it is not known whether the high concentrations of 15d-PGJ<sub>2</sub> required for these responses are consistent with physiological levels. A new study (see the related article beginning on page 945) suggests that in vivo 15d-PGJ<sub>2</sub> levels are actually several orders of magnitude below the levels required to induce many of the biological effects attributed to this molecule.

*J. Clin. Invest.* **112**:828-830 (2003). doi:10.1172/JCI200319796.

Over the past several years there has been increasing interest in the prostaglandin D<sub>2</sub>-derived (PGD<sub>2</sub>-derived) product 15-deoxy- $\Delta^{12,14}$ -PGJ<sub>2</sub> (15d-PGJ<sub>2</sub>) (Figure 1). This substance was first identified in 1983 as a degradation product of PGD<sub>2</sub>, formed after incubation for extended periods of time in the presence of albumin (1). 15d-PGJ<sub>2</sub> attracted relatively little attention until 1995, when Forman et al. (2) and Kliewer et al. (3) independently reported in *Cell* that it is capable of activating the transcription factor PPAR $\gamma$ . This finding was of considerable interest because the endogenous ligand for PPAR $\gamma$  was unknown, in

spite of the availability of a number of pharmacological agonists that are used in the treatment of type 2 diabetes because of their ability to enhance sensitivity to insulin. The hypothesis that 15d-PGJ<sub>2</sub> fulfilled this role was thus very attractive, and 15d-PGJ<sub>2</sub> is often referred to as an endogenous PPAR $\gamma$  ligand. For this to be correct, 15d-PGJ<sub>2</sub> would have to act at concentrations compatible with its physiological levels. Bell-Parikh et al. now report a highly sensitive and specific assay that they have used to address this question (4).

### Conversion of PGD<sub>2</sub> to 15d-PGJ<sub>2</sub>

The synthesis of prostaglandins is initiated by activation of phospholipase A<sub>2</sub>, resulting in the release of arachidonic acid, which is then converted by cyclooxygenases to PGH<sub>2</sub>. This unstable intermediate is converted enzymatically to a series of biologically active prostanoids, each of which has its own selective receptor(s). PGD<sub>2</sub> is formed by the actions of hematopoietic and lipocalin-type PGD<sub>2</sub> synthases and is metabolized enzymatically to

11 $\beta$ -PGF<sub>2 $\alpha$</sub>  and 13,14-dihydro-15-keto-PGD<sub>2</sub>. Alternatively, PGD<sub>2</sub> can undergo chemical dehydration, losing a molecule of water to form the cyclopentenone prostaglandin PGJ<sub>2</sub>. PGJ<sub>2</sub> can then undergo further dehydration by loss of the 15-hydroxyl group, which, coupled with migration of the 13,14-double bond of PGJ<sub>2</sub>, results in the formation of 15d-PGJ<sub>2</sub> (Figure 1). These reactions are promoted by albumin but proceed at a relatively slow rate compared to the very rapid formation of PGs from PGH<sub>2</sub> by prostanoid synthases. There is no evidence for the enzymatic formation of any of these cyclopentenone PGs, and this lack of evidence is one of the factors that has brought into question their physiological relevance.


### 15d-PGJ<sub>2</sub> as a PPAR $\gamma$ ligand

The two papers published in *Cell* in 1995 sparked considerable research on the biological effects of 15d-PGJ<sub>2</sub>. As it has been considered to be an endogenous ligand for PPAR $\gamma$ , 15d-PGJ<sub>2</sub> has been employed in many studies on the role of this transcription factor and in most cases shown to have effects similar to those of synthetic PPAR $\gamma$  ligands. This raised the possibility that 15d-PGJ<sub>2</sub> could be a physiological regulator of processes affected by PPAR $\gamma$ , including adipocyte differentiation and regulation of glucose levels (5). In addition, there has recently been considerable interest in the role of PPAR $\gamma$  in regulating the inflammatory response, as 15d-PGJ<sub>2</sub> and other PPAR $\gamma$  agonists inhibit the expression of a variety of proteins with proinflammatory properties, including cyclooxygenase-2, iNOS, and a variety of cytokines (6). These effects appear to be mediated by inhibitory effects on various transcription factors, including NF- $\kappa$ B, activator protein-1 (AP-1), and signal transducer and activator of transcription-1 (STAT-1). This has led

**Address correspondence to:** William S. Powell, Meakins-Christie Laboratories and Department of Medicine, McGill University, 3626 St. Urbain Street, Montreal, Quebec H2X 2P2, Canada. Phone: (514) 398-3864 ext. 094071; Fax: (514) 398-7483; E-mail: William.Powell@McGill.ca.

**Conflict of interest:** The author has declared that no conflict of interest exists.

**Nonstandard abbreviations used:** prostaglandin D<sub>2</sub> (PGD<sub>2</sub>); 15-deoxy- $\Delta^{12,14}$ -PGJ<sub>2</sub> (15d-PGJ<sub>2</sub>); the concentration of agonist that elicits 50% of the maximal response (EC<sub>50</sub>).



**Figure 1**

Formation of 15d-PGJ<sub>2</sub> and its effects on receptors and signaling molecules. The degradation of PGD<sub>2</sub> to 15d-PGJ<sub>2</sub> is enhanced by albumin. In addition to its effects on intracellular proteins, 15d-PGJ<sub>2</sub> also activates the DP<sub>2</sub> receptor. PGD<sub>2</sub> can also be metabolized enzymatically (Enz) to 11β-PGF<sub>2α</sub> and 13,14-dihydro-15-keto-PGD<sub>2</sub> (dhk-PGD<sub>2</sub>). The latter metabolite also activates the DP<sub>2</sub> receptor. Some of the effects of 15d-PGJ<sub>2</sub> on intracellular proteins are mediated by Michael addition to the reactive cyclopentenone ring (box). In contrast to its effect on the DP<sub>2</sub> receptor, 15d-PGJ<sub>2</sub> does not activate the DP<sub>1</sub> receptor.

to the proposal that 15d-PGJ<sub>2</sub> may be an anti-inflammatory mediator that plays an important role in the resolution of inflammation (7). There is also evidence that PPARγ activators may have antitumorigenic effects due to their inhibitory effects on tumor cell proliferation and angiogenesis.

Although it is clear that 15d-PGJ<sub>2</sub> can stimulate PPARγ, the physiological and pharmacological relevance of this may be debatable for several reasons. First, the concentrations required to activate PPARγ are generally reported to be in the μM range, which is very high for prostaglandins, since they usually act at low nM concentrations and are normally formed only in small amounts. However, 15d-PGJ<sub>2</sub> acts intracellularly, in contrast to other PGs, which act principally as extracellular mediators, interacting with receptors on the cell surface. Thus it may be argued that it is difficult to know the intracellular concentration of 15d-PGJ<sub>2</sub>. These key issues are addressed by Bell-Parikh et al. (4). Secondly, there is growing evidence that 15d-PGJ<sub>2</sub> can induce a variety of responses independently of PPARγ, some of which would oppose those expected by activation of PPARγ. This

lack of selectivity could limit the potential pharmacological usefulness of this substance.

#### Other effects of 15d-PGJ<sub>2</sub>

The expression of PPARγ is more restricted than is the expression of other members of the PPAR family (8), and 15d-PGJ<sub>2</sub> has been shown to induce responses in cells devoid of this receptor (9). Some of these effects may be mediated through the covalent binding of 15d-PGJ<sub>2</sub> to proteins. This is due to the reactive cyclopentenone ring of 15d-PGJ<sub>2</sub>, which readily reacts with substances containing nucleophilic groups such as cysteinyl thiol groups of proteins. Such reactions are termed Michael addition reactions and also occur with 2-cyclohexenone itself. Thus 15d-PGJ<sub>2</sub>, in addition to inhibiting NF-κB activation via PPARγ, has been shown to covalently bind to IκB kinase, inhibiting its activity and causing NF-κB to remain in an inactive state. 15d-PGJ<sub>2</sub> also inhibits the binding of NF-κB subunits to DNA by a similar mechanism (10). Similarly, 15d-PGJ<sub>2</sub> was found to form an adduct with cysteine-184 of H-Ras in 3T3 cells, but not with either N-Ras or K-Ras, which do not contain a cysteine

residue in this position (11). However, in contrast to its inhibitory effect on NF-κB, 15d-PGJ<sub>2</sub> activates H-Ras, resulting in the stimulation of MAPK and PI-3K, which appear to be responsible for the positive effect of 15d-PGJ<sub>2</sub> on cell proliferation.

In addition to the intracellular effects of 15d-PGJ<sub>2</sub> on transcription factors and signaling molecules, 15d-PGJ<sub>2</sub> can also act on at least one cell membrane receptor. The parent compound, PGD<sub>2</sub>, acts through two such receptors: the DP<sub>1</sub> receptor and the DP<sub>2</sub> receptor, which is also known as the chemoattractant receptor-homologous molecule expressed on TH2 cells (CRTH2). We have shown that this substance activates the DP<sub>2</sub> receptor on eosinophils with a potency (concentration of agonist that elicits 50% of the maximal response [EC<sub>50</sub>] ~10 nM) nearly equal to that of PGD<sub>2</sub>, the main ligand for this receptor (12). This effect thus occurs at a much lower concentration than that required for activation of PPARγ and would be pro-inflammatory rather than anti-inflammatory.

#### Is 15d-PGJ<sub>2</sub> biologically relevant?

15d-PGJ<sub>2</sub> clearly induces or inhibits a variety of PPARγ-dependent and PPARγ-independent responses and is active both in vitro and in vivo. However, the high concentrations required to induce of these effects brings into question their physiological relevance, as PGs normally exhibit EC<sub>50</sub> values in the low nM range. Nevertheless, the number of publications on 15d-PGJ<sub>2</sub> in high-profile journals continues to increase rapidly despite the paucity of information on its concentrations in biological fluids. Measurement of 15d-PGJ<sub>2</sub> in a physiologically relevant setting has been hampered by the lack of reliable immunoassay methods. To raise antibodies to 15d-PGJ<sub>2</sub> and other prostaglandins, these small molecules must first be conjugated to proteins. As 15d-PGJ<sub>2</sub> spontaneously forms such conjugates in biological milieu, it is difficult to know how much of the immunoreactive material is free 15d-PGJ<sub>2</sub> and how much is 15d-PGJ<sub>2</sub> or related lipids bound to proteins or glutathione. Because of such uncertainties, one commercially available

immunoassay (Cayman Chemical Co., Ann Arbor, Michigan, USA) has been withdrawn from the market. Another antibody has been reported to detect relatively high levels (~80 nM) of 15d-PGJ<sub>2</sub> in medium from RAW264.7 macrophage-like cells stimulated with LPS (13). However, additional validation of this assay would be desirable. The gold standard for measurement of eicosanoids in biological fluids is clearly mass spectrometry, and the availability of reliable mass spectrometric methods to quantitate 15d-PGJ<sub>2</sub> is essential to accurately estimate its biological levels as well as to validate immunological assays. In this issue of the *JCI*, Bell-Parikh et al. (4) report a highly sensitive liquid chromatography/tandem mass spectrometry assay for this substance. The authors show that although 3T3-L1 fibroblasts can produce a very small amount of 15d-PGJ<sub>2</sub> (~5 pM), its production does not increase during differentiation of these cells to adipocytes. Furthermore, the amounts of 15d-PGJ<sub>2</sub> released into the medium are more than five orders of magnitude lower than the amounts required for PPAR $\gamma$ -dependent differentiation of these cells into adipocytes, and the concentration of cell-associated 15d-PGJ<sub>2</sub> (~1 nM) is over three orders of magnitude lower. The levels of 15d-PGJ<sub>2</sub> excreted in the urine were also found to be very low compared to the levels of other PGs and were not altered in diabetes or following administration of LPS. 15d-PGJ<sub>2</sub> was barely detectable in synovial fluid. These results provide strong evidence that, although 15d-PGJ<sub>2</sub> can be produced in vivo, its levels are far too low to be compatible with a role for this substance as an endogenous activator of PPAR $\gamma$ . Furthermore, because of its

lack of selectivity for PPAR $\gamma$ , 15d-PGJ<sub>2</sub> may be of limited usefulness in defining the role of this receptor.

### Conclusions

The study by Bell-Parikh et al. (4), using a highly sensitive and selective assay, suggests that the amounts of 15d-PGJ<sub>2</sub> generated in vivo are insufficient for it to play a physiological role in modulating most of the responses, including activation of PPAR $\gamma$ , that it has been shown to induce at  $\mu$ M concentrations in vitro. There are only limited instances in which 15d-PGJ<sub>2</sub> has been demonstrated to act at low (1–10) nM levels, such as activation of eosinophils through the DP<sub>2</sub> receptor (12) and stimulation of tumor cell proliferation (14). Further information on the ability of certain cell types such as macrophages (13) to produce 15d-PGJ<sub>2</sub>, using mass spectrometric techniques such as those described in the current issue, would seem important in determining whether it could possibly affect such processes. The usefulness of 15d-PGJ<sub>2</sub> at a pharmacological level is also rather questionable, as it has been shown to elicit a variety of responses, many of them in opposition to one another. For example, it has been reported to be antiangiogenic (15), yet also to induce VEGF synthesis (16), and it both induces and prevents proliferation of tumor cells (17). Because of its reactivity, 15d-PGJ<sub>2</sub>, like cyclopentenone itself, can react with proteins and either inhibit or enhance their activity. Therefore, if used at sufficiently high concentrations, it will certainly induce or inhibit responses. While this in itself may be of interest, it does not necessarily add to our understanding of physiologically relevant regulatory mechanisms.

1. Fitzpatrick, F.A., and Wynalda, M.A. 1983. Albumin-catalyzed metabolism of prostaglandin D<sub>2</sub>. Identification of products formed in vitro. *J. Biol. Chem.* **258**:11713–11718.
2. Forman, B.M., et al. 1995. 15-Deoxy- $\Delta^{12,14}$ -prostaglandin J<sub>2</sub> is a ligand for the adipocyte determination factor PPAR $\gamma$ . *Cell*. **83**:803–812.
3. Kliewer, S.A., et al. 1995. A prostaglandin J<sub>2</sub> metabolite binds peroxisome proliferator-activated receptor- $\gamma$  and promotes adipocyte differentiation. *Cell*. **83**:813–819.
4. Bell-Parikh, L.C., et al. 2003. Biosynthesis of 15-deoxy- $\Delta^{12,14}$ -PGJ<sub>2</sub> and the ligation of PPAR $\gamma$ . *J. Clin. Invest.* **112**:945–955. doi:10.1172/JCI200318012.
5. Murphy, G.J., and Holder, J.C. 2000. PPAR- $\gamma$  agonists: therapeutic role in diabetes, inflammation and cancer. *Trends Pharmacol. Sci.* **21**:469–474.
6. Daynes, R.A., and Jones, D.C. 2002. Emerging roles of PPARs in inflammation and immunity. *Nat. Rev. Immunol.* **2**:748–759.
7. Gilroy, D.W., et al. 1999. Inducible cyclooxygenase may have anti-inflammatory properties. *Nat. Med.* **5**:698–701.
8. Braissant, O., Foulfelle, F., Scotto, C., Dauca, M., and Wahli, W. 1996. Differential expression of peroxisome proliferator-activated receptors (PPARs): tissue distribution of PPAR- $\alpha$ , - $\beta$ , and - $\gamma$  in the adult rat. *Endocrinology*. **137**:354–366.
9. Chawla, A., et al. 2001. PPAR- $\gamma$  dependent and independent effects on macrophage-gene expression in lipid metabolism and inflammation. *Nat. Med.* **7**:48–52.
10. Straus, D.S., et al. 2000. 15-Deoxy- $\Delta^{12,14}$ -prostaglandin J<sub>2</sub> inhibits multiple steps in the NF- $\kappa$ B signaling pathway. *Proc. Natl. Acad. Sci. U. S. A.* **97**:4844–4849.
11. Oliva, J.L., et al. 2003. The cyclopentenone 15-deoxy- $\Delta^{12,14}$ -prostaglandin J<sub>2</sub> binds to and activates H-Ras. *Proc. Natl. Acad. Sci. U. S. A.* **100**:4772–4777.
12. Monneret, G., Li, H., Vasilescu, J., Rokach, J., and Powell, W.S. 2002. 15-Deoxy- $\Delta^{12,14}$ -prostaglandins D<sub>2</sub> and J<sub>2</sub> are potent activators of human eosinophils. *J. Immunol.* **168**:3563–3569.
13. Shibata, T., et al. 2002. 15-deoxy- $\Delta^{12,14}$ -prostaglandin J<sub>2</sub>. A prostaglandin D<sub>2</sub> metabolite generated during inflammatory processes. *J. Biol. Chem.* **277**:10459–10466.
14. Chinery, R., et al. 1999. Prostaglandin J<sub>2</sub> and 15-deoxy- $\Delta^{12,14}$ -prostaglandin J<sub>2</sub> induce proliferation of cyclooxygenase-depleted colorectal cancer cells. *Cancer Res.* **59**:2739–2746.
15. Xin, X., Yang, S., Kowalski, J., and Gerritsen, M.E. 1999. Peroxisome proliferator-activated receptor  $\gamma$  ligands are potent inhibitors of angiogenesis in vitro and in vivo. *J. Biol. Chem.* **274**:9116–9121.
16. Bamba, H., Ota, S., Kato, A., Kawamoto, C., and Fujiwara, K. 2000. Prostaglandins up-regulate vascular endothelial growth factor production through distinct pathways in differentiated U937 cells. *Biochem. Biophys. Res. Commun.* **273**:485–491.
17. Clay, C.E., et al. 2001. Magnitude of peroxisome proliferator-activated receptor- $\gamma$  activation is associated with important and seemingly opposite biological responses in breast cancer cells. *J. Invest. Med.* **49**:413–420.